

전공의 연수강좌, 강촌

양대혈관우심실 기시증

신홍주

세브란스 심장혈관병원 심장혈관외과

May 21 / 2015

Definition of DORV


- A congenital anomaly in which both great arteries arise wholly or in large part from the right ventricle
- Presence of bilateral infundibulum
- Presence of atrioventricular valve-semilunar valve discontinuity (mitral aortic discontinuity)
- In the spectrum of TOF and TGA
- **50% rule**: a heart is termed DORV if > 50% of both great arteries arise from the right ventricle

History of nomenclature


- 1898, Vierordt partial TGA
- 1923, Spitzer type II TGA (simple TGA)
- 1949, Taussig and Bing TGA with levoposition of PA
- 1950, Lev and Bolk Taussig-Bing heart
- 1952, Braun double outlet ventricle
- 1957, Witham DORV

- 1957, Kirklin at Mayo Clinic 1st operation
- 1967, Kirklin successful repair of Taussig-Bing
- 1968, McGoon
- 1971, Kawashima

Classification by Lev (1972)





DORV with subaortic VSD



DORV with subpulmonic VSD

Classification by Lev (1972)



DORV with doubly committed VSD

DORV with noncommitted VSD

CHSS classification

Congenital Heart Surgery Nomenclature and Database Project: Double Outlet Right Ventricle

Henry L. Walters III, MD, Constantine Mavroudis, MD, Christo I. Tchervenkov, MD, Jeffrey P. Jacobs, MD, François Lacour-Gayet, MD, and Marshall L. Jacobs, MD

Departments of Surgery, Wayne State University School of Medicine, Children's Hospital of Michigan, Detroit, Michigan; Northwestern University School of Medicine, Children's Memorial Hospital, Chicago, Illinois; McGill University, The Montreal Children's Hospital, Montreal, Quebec, Canada; University of South Florida School of Medicine, All Children's Hospital, St. Petersburg, Florida; Marie Lannelongue Hospital, Paris, France; and Hanneman University School of Medicine, St. Christopher's Children's Hospital, Philadelphia, Pennsylvania

Double outlet right ventricle (DORV) is a type of ventriculoarterial connection in which both great vessels arise entirely or predominantly from the right ventricle. Although the presence of aortic-mitral discontinuity and bilateral conus are important descriptors, they should not serve as absolute prerequisites for the diagnosis of DORV. The morphology of DORV is encompassed by a careful description of the ventricular septal defect (VSD) with its relationship to the semilunar valves, the great artery relationships to each other, the coronary artery anatomy, the presence or absence of pulmonary outflow tract obstruction (POTO) and aortic outflow tract obstruction (AOTO), the tricuspid-pulmonary annular distance,


and the presence or absence of associated cardiac lesions. The preferred surgical treatment involves the connection of the left ventricle to the systemic circulation by an intraventricular tunnel repair connecting the VSD to the systemic semilunar valve. This ideal surgical therapy is not always possible due to the presence of confounding anatomical barriers. A multitude of alternative surgical procedures has been devised to accommodate these more complex situations. A framework for the development of the DORV module for a pediatric cardiac surgical database is proposed.


(Ann Thorac Surg 2000;69:S249-63)



© 2000 by The Society of Thoracic Surgeons


CHSS classification of DORV

- DORV, VSD type
 - Subaortic or doubly committed, without RVOTO
- DORV, TOF type
 - Subaortic or doubly committed, with RVOTO
- DORV, TGA type
 - Subpulmonary without PS (**Taussig-Bing**) or with PS
- DORV, Remote type
 - Non committed, with or without RVOTO
- DORV, IVS













Normal




TOF



DORV, Subaortic



DORV, Subpulmonary



Complete Transposition

Segments of Muscular Septum



Anterior Posterior

Insertion of Conus Septum



Parietal Septal

Bulboventricular Foramen



Atrioventricular Semilunar Continuity



Figure 23-1, Diagrammatic representations of transverse sections of specimens that represent the spectrum of developmental abnormalities in hearts with abnormal ventriculoarterial connection. Ao, aorta; PA, pulmonary artery




Subaortic VSD

| <i>n</i> = 73 | SubAo VSD <i>n</i> (%) | SubP VSD <i>n</i> (%) |
|---------------------|------------------------------|-----------------------------|
| | 31(42) | 27(37) |
| Pulmonary stenosis | 15(48) | 5(19) |
| Aortic stenosis | 4(13) | 4(15) |
| Ao arch obstruction | 1(3) | 14(52) |
| TPD < Ao annulus | 2(6) | 14(52) |

TPD = Tricuspid to pulmonary valve distance



Subpulmonary VSD



Noncommitted
VSD

| <i>n</i> = 73 | Noncommitted <i>n</i> (%) | Doubly committed <i>n</i> (%) |
|---------------------|------------------------------|----------------------------------|
| | 10(14) | 5(7) |
| Pulmonary stenosis | 5(50) | 2(40) |
| Aortic stenosis | 3(30) | 3(60) |
| Ao arch obstruction | 2(20) | 2(40) |
| TPD < Ao annulus | 4(40) | 0 |

TPD = Tricuspid to pulmonary valve distance



Doubly committed
VSD

Figure 23.9 Important anatomic features of VSD groups in a review of 73 patients who underwent biventricular repair of DORV between 1981 and 1991 at Children's Hospital Boston. (From Aoki et al. Results of biventricular repair for double outlet right ventricle. *J Thorac Cardiovasc Surg*, 1994; 107:340, with permission from Elsevier.)




Fig. 24-3 The relationship of the VSD to the great arteries in DORV. **A**, Subaortic VSD without pulmonary stenosis. **B**, Subaortic VSD with pulmonary stenosis. **C**, Subpulmonary VSD (Taussig-Bing malformation). **D**, Doubly-committed VSD. **E**, Noncommitted (remote) VSD. **F**, Intact interventricular septum.




Figure 32.2




Figure 32.3




Figure 32.4




Figure 32.6




Figure 32.7

TV-PV distance





Figure 32.8






Figure 32.9




Figure 32.11




Figure 32.12




Figure 23-3. Kawashima Intraventricular repair: Artist's cut-away view of the Taussig-Bing heart with side-by-side great arteries, (A) The subpulmonic ventricular septal defect (VSD) and subaortic conus are represented in relationship to the great arteries and semilunar valves, (B) The excised subaortic conus is shown in preparation for (C) the left ventricular-to-aorta Intraventricular tunnel, shown here constructed with a Dacron patch and interrupted pledged sutures.






Figure 23-4. Arterial switch, ventricular septal defect closure, and aortic relocation: one-stage neonatal repair of (A) the Taussig-Bing heart with subaortic obstruction, small ascending aorta, and coarctation of the aorta, (B) After great artery transection and coronary artery transfer are accomplished, the coarctation is repaired under circulatory arrest by coarctectomy with end-to-end anastomosis between the “proximal” ascending and descending aorta. A longitudinal incision is made in the lateral ascending aorta, which is now oriented in a transverse position for anastomosis to the neo-aorta, (C) Completed repair showing the neopulmonary artery reconstruction with a valved homograft after subvalvular resection.



(a)



(b)

Figure 23.2 (a) Double outlet right ventricle with subaortic VSD. Physiologically this anomaly is similar to tetralogy of Fallot. Left ventricular blood is predominantly directed to the aorta after passing through a subaortic VSD. (b) Double outlet right ventricle with subpulmonary VSD. Physiologically this anomaly is similar to transposition of the great arteries. Left ventricular blood passes through the subpulmonary VSD into the pulmonary arteries so that oxygen saturation is higher in the pulmonary artery than in the aorta.




Figure 23.4 (a) A prominent conal septum may project into the baffle pathway; (b) Resection of the conal septum may be necessary to prevent subaortic stenosis within the baffle pathway; (c) Completion of intraventricular repair with a baffle.







Figure 23.5 (a) Intraventricular repair of double outlet right ventricle with a subaortic VSD that is moderately distant from the aortic annulus. (b) Autologous pericardium is preferred for baffle construction (or 'stretch' PTFE). Both these materials are less likely to acquire a thick fibrous pseudointima than Dacron. There is also greater pliability of the baffle so that a kink is less likely to project into the central point of the pathway. (c) Beyond infancy the baffle pathway may be so long that it is impractical to use interrupted pledgetted sutures. A running technique is a reasonable alternative. (d) The completed left ventricle to aorta baffle pathway as part of an intraventricular repair of DORV.



(a)



(b)



(c)

Figure 23.6 Rastelli repair for DORV. (a) This patient is close to the transposition end of the DORV spectrum so that the pulmonary valve is almost in fibrous continuity with the mitral valve. (b) The main pulmonary artery has been divided, the pulmonary valve has been excised and the stump of the proximal main pulmonary artery has been oversewn. A pericardial or stretch PTFE baffle is being sutured into place to direct left ventricular blood through the VSD to the aorta. The former pulmonary outflow is incorporated under the baffle. (c) The Rastelli repair is completed by placing a homograft conduit between the right ventricle and distal divided main pulmonary artery.









Figure 53-15 Repair of double outlet right ventricle with subpulmonary ventricular septal defect (VSD) (Taussig-Bing heart) when there is only a short distance between tricuspid and pulmonary valves (Patrick-McGoon method^{MS,PT,P3}). Aorta is usually anterior to pulmonary trunk. **A**, Through a transverse right ventriculotomy, the VSD is enlarged and much of the inlet septum excised (*dashed line*). **B**, A contoured polyester (or polytetrafluoroethylene) patch is cut from a tube graft whose diameter is about 20% larger than that of the aorta. Initial suturing is performed in the usual manner, preventing damage to the bundle of His. A continuous suture is used, often supplemented by interrupted stitches. **C**, With the leftward arm of the continuous suture, suture line is continued leftward along the posterior and then leftward margin of the ventriculopulmonary trunk junction, and finally along the anterior margin of this junction (*dashed line*). With the other arm of the suture, the patch is sutured inferior to the VSD, to the right side of the septum, and then anterior to the defect. **D**, Insertion of spiraled patch has been completed. Left ventricular blood now passes beneath the patch to the aorta, while right ventricular blood passes to the pulmonary trunk behind the tunnel.




Surgery and timing for repair

| | | |
|------------|--|--|
| VSD type | About 3 months | Intraventricular repair (IVR) |
| TOF type | 3 - 6 month | IVR + RVOT rec. |
| TGA type | Early and late infancy | <ul style="list-style-type: none"> •ASO + VSD tunnel •Senning/Mustard + VSD tunnel •DKS+VSD tunnel+RV-PA •REV •Kawashima •Nikaidoh |
| Remote VSD | <p>Infancy?</p> <p>After palliation?</p> | <ul style="list-style-type: none"> •Biventricular repair •Univentricular repair |

Intraventricular repair




Conal resection




VSD extension

DORV with subaortic VSD, PS

- Valved conduit
 - Major coronary arteries crossing RV
 - High PVR or distal pulmonary obstruction
- Transannular patch



DORV, subpulmonic VSD (Taussig-Bing Anomaly)




DORV, subpulmonic VSD (Taussig-Bing Anomaly)

Taussig-Bing 기형에서의 대혈관치환술은 완전대혈관전 위증과 비교하여 몇 가지 차이점이 있다.


- 1) 관상동맥 전이 방법의 차이이다. 완전대혈관전위증의 경우 통상적인 관상동맥 배열이 흔한 반면, Taussig-Bing 기형의 경우 다양하고 복잡한 형태의 관상동맥 배열 (single coronary artery, intramural coronary artery) 이 관찰된다
- 2) 둘째, Taussig-Bing 기형에서 대혈관의 배열이 다양한 데, 측측 배열의 대혈관에서 Lecompte 수기(maneuver)의 시행 여부이다
- 3) Taussig-Bing 기형에서 대동맥과 폐동맥의 심한 크기 차이를 보인다 상행대동맥이 폐동맥에 비해 심하게 작으며, 특히 대동맥궁협착이 있는 경우 차이는 더 심해진다
- 4) 대동맥하 협착, 대동맥 축착의 동반이 많다

Damus-Kaye-Stansel procedure, tunnel closure of VSD and RV-PA conduit


- Significant subaortic stenosis
- Too risky to perform arterial switch (intramural coronary artery, single coronary artery)
- Aortic valve insufficiency - aortic valve should be closed



DORV with noncommitted VSD



Intraventricular tunneling



Intraventricular conduit

DORV with noncommitted VSD

- Uni-ventricular repair

Anomalies of the atrioventricular valves


Multiple muscular VSDs

Inability to reliably channel the remote

VSD to aorta

- Hypoplasia of the right or left ventricle

DORV decision tree



DORV decision tree

Complex DORV /TGA type

Aorta anterior,
no PS

Aorta anterior,
PS

Side by Side
GVs, no PS

Arterial switch

Rastelli/Nikaidoh

Normal AoVA,
adequate TV-
PV diam

Small AoVA arch
obstruction

IVR (tunnel)

Complex Arterial
switch

Complication

- Heart block
- Residual left-to-right shunt
- **LVOT obstruction**
 - Inadequate enlargement of VSD
 - Poor configuration of intraventricular tunnel patch
- **RVOT obstruction**
 - Obstructing muscle bundle or patch
- Myocardial dysfunction
- Coronary ischemia

Table 23-1. Summary of surgical treatment of remote DORV

| Authors (year) | Time period of operations | No of patients | Previous palliations | Techniques of repair | Age at repair | Early death | Follow-up | Late death | Reoperation |
|--|---------------------------|----------------|---|---------------------------|---------------------|-------------|---------------------|------------|--|
| Belli (1999) ^[41] | 1987-1997 | 23 | 9 PAB (7) COA repair (5) BTS (2) | IVR 21 ASO 2 | 20no (50d-10yr) | 2 (9%) | 58mo | No | 8 (35%) Subaortic stenosis (6) TV repair + RVOTR(1) AVR (1) |
| Barbero-Marcial (1999) ^[43] | 1987-1999 | 18 | 7 BTS (4) PAB (3) | IVR (Multiple patches) | 4.7yr (2mo-13yr) | 2 (11%) | 2.7yr (1mo-12yr) | 3 (16.6%) | MV repair (1) |
| Lacour-Gayet (2002) ^[77] | 1998-2001 | 10 | 7 PAB | ASO | 16mo (3wk-4.5yr) | 1 | 20mo (4mo-4yr) | No | No |
| Artrip (2006) ^[21] | 2000-2005 | 10 | 8 BTS (3) PAB (5) COA repair (2) | IVR 7 ASO 3 | 11mo (9d-4yr) | 1 | 20mo | 1 | NA |
| Hu (2010) ^[79] | 2006-2009 | 6 | NA | Double-root translocation | *4yr (1-16yr) | No | *22mo (2-36mo) | 없음 | No |

(ASO; arterial switch operation after LV-PA tunneling, AVR; aortic valve replacement, BTS; Blalock-Taussig shunt, COA; coarctation of the aorta, IVR; intraventricular rerouting, MV; mitral valve, PAB; pulmonary artery banding, TV; tricuspid valve)

NA; not available

* 4 patients with Taussig-Bing anomaly included

References

- 김용진, 심장외과학
23 chapter DORV by
이창하
- Stark J, et al. Surgery
for congenital heart
defects, 3rd edition.
John Wiley & Sons,
Ltd
- Kirklin / Barratt-Boyes.
Cardiac surgery, 4th
edition. ELSEVIER
- Jonas RA.
Comprehensive
surgical management
of congenital heart
disease, 1st edition,
2004. Arnold, part of
Hodder Education

SURGERY for CONGENITAL HEART DEFECTS

Third Edition

Editors

J STARK, M. de LEVAL and VT TSANG

*Great Ormond Street Hospital
for Children NHS Trust, London*


Illustrated by

MICHAEL COURTNEY



John Wiley & Sons, Ltd

COMPREHENSIVE SURGICAL MANAGEMENT OF CONGENITAL HEART DISEASE



RICHARD A. JONAS

ILLUSTRATED BY REBEKAH DODSON



Get Full Access and More at

ExpertConsult.com

Volume 1

Kirclin/Barratt-Boyes

Fourth Edition

Cardiac Surgery

NICHOLAS T. KOUCHOUKOS • EUGENE H. BLACKSTONE
FRANK L. HANLEY • JAMES K. KIRKLIN

ELSEVIER
www.elsevier.com